

## SCIENTIFIC NOTE

**FIRST RECORD OF *DASYCORIXA RAWSONI*  
(HEMIPTERA: CORIXIDAE) IN THE UNITED STATES<sup>1</sup>**Bruce A. Hanson<sup>2</sup>, Ned H. Euliss, Jr.<sup>2</sup>, David M. Mushet<sup>2</sup>, and Steve W. Chordas III<sup>3</sup>

Hungerford (1948) described the genus *Dasycorixa* in his monograph of the Corixidae of the Western Hemisphere. This genus contains the three species *Dasycorixa hybrida* (Hungerford 1926), *Dasycorixa johanseni* (Walley, 1931), and *Dasycorixa rawsoni* (Hungerford 1948), all of which are known from Canada. Prior to this paper, only one *Dasycorixa* species (*D. hybrida*) was known from the United States. The purpose of this note is to report an additional *Dasycorixa* species (*D. rawsoni*, Fig. 1) as a new country record for the United States.

Types of *Dasycorixa rawsoni* collected from Lizard Lake in Saskatchewan, Canada, on August 10, 1938, were archived in the Francis Huntington Snow Entomological collection, University of Kansas. This species has been collected in Alberta, British Columbia, Manitoba, and the Northwest Territories.




Fig. 1. Dorsal view of *Dasycorixa rawsoni* (Hungerford 1948).

<sup>1</sup> Received on August 16, 2002. Accepted on March 13, 2004.

<sup>2</sup> U. S. Geological Survey, Northern Prairie Wildlife Research Center, 8711 37th Street SE, Jamestown, North Dakota 58401, U.S.A. E-mail of author BAH: bruce\_hanson@usgs.gov.

<sup>3</sup> The Ohio State University, Museum of Biological Diversity, 1315 Kinnear Road, Columbus, Ohio 43212, U.S.A.

In July 1999, we collected two males of *D. rawsoni* at the Cottonwood Lake Study Area. This area is located in Stutsman County, North Dakota, and was described by Swanson (1978). Voucher specimens are archived in the aquatic invertebrate collection at the U.S. Geological Survey, Northern Prairie Wildlife Research Center in Jamestown, North Dakota. Both specimens were captured using a funnel trap (Swanson 1978) set in 84 cm deep water in the deep marsh zone (open water phase) of a semipermanent wetland (Stewart and Kantrud 1971). At the time of sampling, water temperature was 26.0 C, and the specific conductance was 2250  $\mu\text{S cm}^{-1}$ . Aquatic plants in the vicinity of the capture site included sago pondweed [*Stuckenia pectinatus* (L.) Boerner], shortspike water milfoil (*Myriophyllum sibiricum* Komarov), coontail (*Ceratophyllum demersum* L.), star duckweed (*Lemna trisulca* L.), and broad-leaved cattail (*Typha latifolia* L.). As is typical for the prairie pothole region, this wetland fluctuates between wet and dry phases, but it was in lake phase (van der Valk and Davis 1978) and had a maximum water depth of 3.55 m when we collected the two specimens of *D. rawsoni*. The hydrologic setting, geology, water chemistry, and wetland plant communities of focal wetlands at the Cottonwood Lake Study Area have been described by Winter and Carr (1980), Swanson (1990), LaBaugh et al. (1996), and Poiani et al. (1996), respectively.

#### ACKNOWLEDGMENTS

We thank Jane E. Austin, Lawrence D. Igl, and Dale A. Wrubleski for reviewing the manuscript.

#### LITERATURE CITED

- Hungerford, H. B. 1948. The Corixidae of the Western Hemisphere. University of Kansas Science Bulletin 32:827.
- LaBaugh, J. W., T. C. Winter, G. A. Swanson, D. O. Rosenberry, R. D. Nelson, and N. H. Euliss Jr. 1996. Changes in atmospheric patterns affect midcontinent wetlands sensitive to climate. *Limnology and Oceanography* 41:864-870.
- Poiani, K. A., W. C. Johnson, G. A. Swanson, and T. C. Winter. 1996. Climate change and northern prairie wetlands. *Limnology and Oceanography* 41:871-881.
- Stewart, R. E. and H. A. Kantrud. 1971. Classification of natural ponds and lakes in the glaciated prairie region. U.S. Fish and Wildlife Service, Resource Publication No. 92. 57 pp.
- Swanson, G. A. 1978. Funnel trap for collecting littoral aquatic invertebrates. *Progressive Fish-Culturist* 40:73.
- Swanson, G. A. 1987. An introduction to the Cottonwood Lake area. *Proceedings of the North Dakota Academy of Science* 41:25.
- Swanson, K. D. 1990. Chemical evolution of ground water in clay till in a prairie wetlands setting in the Cottonwood Lake area, Stutsman County, North Dakota. M.S. Thesis. University Wisconsin, Madison. 229 pp.
- van der Valk, A. G. and C. B. Davis. 1978. The role of seed banks in the vegetation dynamics of prairie glacial marshes. *Ecology* 59:322-335.
- Winter, T. C. and M. R. Carr. 1980. Hydrologic setting of wetlands in the Cottonwood Lake area, Stutsman County, North Dakota. U.S. Geological Survey. Water Resources Investigation 8-99.